

University of Groningen

Complete Chiral Resolution Using Additive-Induced Crystal Size Bifurcation During Grinding

Noorduyn, Wim L.; Asdonk, Pim van der; Meekes, Hugo; Enckevort, Willem J.P. van; Kaptein, Bernard; Leeman, Michel; Kellogg, Richard M.; Vlieg, Elias

Published in:
Angewandte Chemie International Edition

DOI:
[10.1002/anie.200806214](https://doi.org/10.1002/anie.200806214)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Noorduyn, W. L., Asdonk, P. V. D., Meekes, H., Enckevort, W. J. P. V., Kaptein, B., Leeman, M., Kellogg, R. M., & Vlieg, E. (2009). Complete Chiral Resolution Using Additive-Induced Crystal Size Bifurcation During Grinding. *Angewandte Chemie International Edition*, 48(18), 3278-3280.
<https://doi.org/10.1002/anie.200806214>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Supporting Information

© Wiley-VCH 2009

69451 Weinheim, Germany


Supporting Information

Complete Chiral Resolution Using Additive Induced Crystal Size Bifurcation Under Near-Equilibrium Conditions

*Wim L. Noorduin, Pim van der Asdonk, Hugo Meekes, Willem J. P. van Enkevort, Bernard Kaptein, Michel Leeman, Richard M. Kellogg, Elias Vlieg**

The continuous process of growth and ablation results in a steady-state for the crystal size distribution (CSD). In case an additive stereoselectively hampers the growth, the CSD shifts towards smaller sizes for that enantiomer. Therefore one might expect a distribution showing two separate maxima, one corresponding to the large crystals of the unhampered enantiomer and the other to the many small crystals of the hampered enantiomer.

We examined the ground residues using Scanning Electron Microscopy (SEM), optical microscopy and Focused Beam Reflectance Measurements (SI Figure1). These measurements, however, showed no convincing sign of two separated size distributions for the enantiomers, but rather a uniform distribution centered on a small crystal size.



SI Figure 1. SEM images of racemic **1** ground in the presence of the enantiopure additive **2**. The scale bars are 60 μm (left image) and 20 μm (right image).